arXiv:1208.1641v1 [math.CA] 8 Aug 2012

ON GENERALIZATION OF DIFFERENT TYPE INEQUALITIES FOR SOME CONVEX FUNCTIONS VIA FRACTIONAL INTEGRALS

İMDAT İŞCAN

Abstract. New identity for fractional integrals have been defined. By using of this identity, we obtained new estimates on generalization of Hadamard, Ostrowski and Simpson type inequalities for $s-$convex, quasi-convex, $m-$convex functions via Riemann Liouville fractional integral.

1. Introduction

Following inequalities are well known in the literature as Hermite-Hadamard inequality, Ostrowski inequality and Simpson inequality respectively:

Theorem 1. *Let $f:I\subseteq\mathbb{R}\rightarrow\mathbb{R}$ be a convex function defined on the interval I of real numbers and $a,b\in I$ with $a<b$. The following double inequality holds*

$$f\left(\frac{a+b}{2}\right)\leq\frac{1}{b-a}\int\limits_a^b f(x)dx\leq\frac{f(a)+f(b)}{2}. \tag{1.1}$$

Theorem 2. *Let $f:I\subseteq\mathbb{R}\rightarrow\mathbb{R}$ be a mapping differentiable in I°, the interior of I, and let $a,b\in I^{\circ}$ with $a<b$. If $|f'(x)|\leq M$, $x\in[a,b]$, then we the following inequality holds*

$$\left|f(x)-\frac{1}{b-a}\int\limits_a^b f(t)dt\right|\leq\frac{M}{b-a}\left[\frac{(x-a)^2+(b-x)^2}{2}\right]$$

for all $x\in[a,b]$. The constant $\frac{1}{4}$ is the best possible in the sense that it cannot be replaced by a smaller one.

Theorem 3. *Let $f:[a,b]\rightarrow\mathbb{R}$ be a four times continuously differentiable mapping on (a,b) and $\left\|f^{(4)}\right\|_{\infty}=\sup\limits_{x\in(a,b)}\left|f^{(4)}(x)\right|<\infty$. Then the following inequality holds:*

$$\left|\frac{1}{3}\left[\frac{f(a)+f(b)}{2}+2f\left(\frac{a+b}{2}\right)\right]-\frac{1}{b-a}\int\limits_a^b f(x)dx\right|\leq\frac{1}{2880}\left\|f^{(4)}\right\|_{\infty}(b-a)^2.$$

In [5], Hudzik and Maligranda considered among others the class of functions which are s-convex in the second sense.

2000 *Mathematics Subject Classification.* 26A51, 26A33, 26D10, 26D15.

Key words and phrases. Hermite–Hadamard inequality, Riemann–Liouville fractional integral, Ostrowski inequality, Simpson type inequalities, $s-$convex function, quasi-convex function, $m-$convex function.

Definition 1. *A function* $f:[0,\infty)\to\mathbb{R}$ *is said to be* $s-$*convex in the second sense if*

$$f(\alpha x+\beta y)\leq \alpha^{s}f(x)+\beta^{s}f(y)$$

for all $x,y\in[0,\infty)$, $\alpha,\beta\geq 0$ *with* $\alpha+\beta=1$ *and for some fixed* $s\in(0,1]$. *This class of* $s-$*convex functions in the second sense is usually denoted by* K_s^2.

It can be easily seen that for s = 1, s-convexity reduces to ordinary convexity of functions defined on $[0,\infty)$.

In [13], G. Toader considered the class of m-convexfunctions: another intermediate between the usual convexity and starshaped convexity.

Definition 2. *The function* $f:[0,b]\to\mathbb{R}, b>0$, *is said to be* $m-$*convex, where* $m\in[0,1]$, *if we have*

$$f(tx+m(1-t)y)\leq tf(x)+m(1-t)f(y)$$

for all $x,y\in[0,b]$ *and* $t\in[0,1]$. *We say that* f *is* $m-$*concave if* $(-f)$ *is* $m-$*convex.*

In [10], Pecaric et al. defined quasi-convex functions as following

Definition 3. *A function* $f:[a,b]\to\mathbb{R}$ *is said quasi-convex on* $[a,b]$ *if*

$$f\left(\lambda x+(1-\lambda)y\right)\leq \sup\left\{f(x),f(y)\right\},$$

for any $x,y\in[a,b]$ *and* $\lambda\in[0,1]$.

The notion of quasi-convex functions generalizes the notion of convex functions. Clearly, any convex function is a quasi-convex function. Furthermore, there exist quasi-convex functions which are not convex (see [6]).

We give some necessary definitions and mathematical preliminaries of fractional calculus theory which are used throughout this paper.

Definition 4. *Let* $f\in L[a,b]$. *The Riemann-Liouville integrals* $J_{a^+}^{\alpha}f$ *and* $J_{b^-}^{\alpha}f$ *of oder* $\alpha>0$ *with* $a\geq 0$ *are defined by*

$$J_{a^+}^{\alpha}f(x)=\frac{1}{\Gamma(\alpha)}\int_{a}^{x}(x-t)^{\alpha-1}f(t)dt,\ x>a$$

and

$$J_{b^-}^{\alpha}f(x)=\frac{1}{\Gamma(\alpha)}\int_{x}^{b}(t-x)^{\alpha-1}f(t)dt,\ x<b$$

respectively, where $\Gamma(\alpha)$ *is the Gamma function defined by* $\Gamma(\alpha)=\int_{0}^{\infty}e^{-t}t^{\alpha-1}dt$ *and* $J_{a^+}^{0}f(x)=J_{b^-}^{0}f(x)=f(x)$.

In the case of $\alpha=1$, the fractional integral reduces to the classical integral. Properties concerning this operator can be found [4, 7, 9].

For some recent result connected with fractional integral see [3, 8, 11, 12].

2. GENERALIZED INTEGRAL INEQUALITIES FOR SOME CONVEX FUNCTIONS VIA FRACTIONAL INTEGRALS

Let $f : I \subseteq \mathbb{R} \to \mathbb{R}$ be a differentiable function on I°, the interior of I, throughout this section we will take

$$\begin{aligned} S_f\left(x,\lambda,\alpha,a,b\right) &= (1-\lambda)\left[\frac{(x-a)^{\alpha}+(b-x)^{\alpha}}{b-a}\right]f(x)+\lambda\left[\frac{(x-a)^{\alpha}f(a)+(b-x)^{\alpha}f(b)}{b-a}\right] \\ &\quad -\frac{\Gamma\left(\alpha+1\right)}{b-a}\left[J_{x^-}^{\alpha}f(a)+J_{x^+}^{\alpha}f(b)\right] \end{aligned}$$

where $a,b \in I$ with $a<b$, $x\in[a,b]$, $\lambda\in[0,1]$, $\alpha>0$ and Γ is Euler Gamma function. In order to prove our main results we need the following identity.

Lemma 1. *Let $f : I \subseteq \mathbb{R} \to \mathbb{R}$ be a differentiable function on I° such that $f' \in L[a,b]$, where $a,b\in I$ with $a<b$. Then for all $x\in[a,b]$, $\lambda\in[0,1]$ and $\alpha>0$ we have:*

$$S_f\left(x,\lambda,\alpha,a,b\right)=\frac{(x-a)^{\alpha+1}}{b-a}\int_0^1\left(t^{\alpha}-\lambda\right)f'\left(tx+(1-t)a\right)dt \tag{2.1}$$

$$+\frac{(b-x)^{\alpha+1}}{b-a}\int_0^1\left(\lambda-t^{\alpha}\right)f'\left(tx+(1-t)b\right)dt.$$

Proof. By integration by parts and changing the variable, we can state

$$\int_0^1\left(t^{\alpha}-\lambda\right)f'\left(tx+(1-t)a\right)dt \tag{2.2}$$

$$\begin{aligned} &= \left(t^{\alpha}-\lambda\right)\frac{f\left(tx+(1-t)a\right)}{x-a}\Bigg|_0^1-\int_0^1\alpha t^{\alpha-1}\frac{f\left(tx+(1-t)a\right)}{x-a}dt \\ &= (1-\lambda)\frac{f(x)}{x-a}+\lambda\frac{f(a)}{x-a}-\frac{\alpha}{x-a}\int_a^x\left(\frac{u-a}{x-a}\right)^{\alpha-1}\frac{f(u)}{x-a}du \\ &= (1-\lambda)\frac{f(x)}{x-a}+\lambda\frac{f(a)}{x-a}-\frac{\Gamma\left(\alpha+1\right)}{(x-a)^{\alpha+1}}J_{x^-}^{\alpha}f(a) \end{aligned}$$

and similarly we get

$$\int_0^1\left(\lambda-t^{\alpha}\right)f'\left(tx+(1-t)b\right)dt \tag{2.3}$$

$$\begin{aligned} &= \left(\lambda-t^{\alpha}\right)\frac{f\left(tx+(1-t)b\right)}{x-b}\Bigg|_0^1-\int_0^1\alpha t^{\alpha-1}\frac{f\left(tx+(1-t)b\right)}{x-b}dt \\ &= (1-\lambda)\frac{f(x)}{b-x}+\lambda\frac{f(b)}{b-x}-\frac{\alpha}{b-x}\int_x^b\left(\frac{b-u}{b-x}\right)^{\alpha-1}\frac{f(u)}{b-x}du \\ &= (1-\lambda)\frac{f(x)}{b-x}+\lambda\frac{f(b)}{b-x}-\frac{\Gamma\left(\alpha+1\right)}{(b-x)^{\alpha+1}}J_{x^+}^{\alpha}f(b) \end{aligned}$$

Multiplying both sides of (2.2) and (2.3) by $\frac{(x-a)^{\alpha+1}}{b-a}$ and $\frac{(b-x)^{\alpha+1}}{b-a}$, respectively, and adding the resulting identities we obtain the desired result. ☐

2.1. **For $s-$convex functions.**

Theorem 4. *Let $f : I \subset [0,\infty) \to \mathbb{R}$ be a differentiable function on I° such that $f' \in L[a,b]$, where $a,b \in I^{\circ}$ with $a<b$. If $|f'|^q$ is $s-$convex on $[a,b]$ for some fixed $q \geq 1$, $x \in [a,b]$, $\lambda \in [0,1]$ then the following inequality for fractional integrals holds*

$$\begin{aligned}
& \left|S_f\left(x,\lambda,\alpha,a,b\right)\right| \\
(2.4) \quad \leq \; & A_1^{1-\frac{1}{q}}\left(\alpha,\lambda\right)\left\{\frac{(x-a)^{\alpha+1}}{b-a}\left(\left|f'\left(x\right)\right|^q A_2\left(\alpha,\lambda\right)+\left|f'\left(a\right)\right|^q A_3\left(\alpha,\lambda\right)\right)^{\frac{1}{q}}\right. \\
& \left. +\frac{(b-x)^{\alpha+1}}{b-a}\left(\left|f'\left(x\right)\right|^q A_2\left(\alpha,\lambda\right)+\left|f'\left(b\right)\right|^q A_3\left(\alpha,\lambda\right)\right)^{\frac{1}{q}}\right\}
\end{aligned}$$

where

$$\begin{aligned}
A_1\left(\alpha,\lambda\right) &= \frac{2\alpha\lambda^{1+\frac{1}{\alpha}}+1}{\alpha+1}-\lambda, \\
A_2\left(\alpha,\lambda,s\right) &= \frac{2\alpha\lambda^{1+\frac{s+1}{\alpha}}+s+1}{(s+1)\left(\alpha+s+1\right)}-\frac{\lambda}{s+1}, \\
A_3\left(\alpha,\lambda,s\right) &= \lambda\left[\frac{1-2\left(1-\lambda^{\frac{1}{\alpha}}\right)^{s+1}}{s+1}\right]+\beta\left(\alpha+1,s+1\right)-2\beta\left(\lambda^{\frac{1}{\alpha}};\alpha+1,s+1\right),
\end{aligned}$$

and β is Euler Beta function defined by

$$\beta\left(x,y\right)=\frac{\Gamma(x)\Gamma(y)}{\Gamma(x+y)}=\int_0^1 t^{x-1}\left(1-t\right)^{y-1}dt, \quad x,y>0.$$

Proof. From Lemma 1, property of the modulus and using the power-mean inequality we have

$$\begin{aligned}
& \left|S_f\left(x,\lambda,\alpha,a,b\right)\right| \leq \frac{(x-a)^{\alpha+1}}{b-a}\int_0^1\left|t^{\alpha}-\lambda\right|\left|f'\left(tx+(1-t)\,a\right)\right|dt \\
& +\frac{(b-x)^{\alpha+1}}{b-a}\int_0^1\left|\lambda-t^{\alpha}\right|\left|f'\left(tx+(1-t)\,b\right)\right|dt \\
\leq \; & \frac{(x-a)^{\alpha+1}}{b-a}\left(\int_0^1\left|t^{\alpha}-\lambda\right|dt\right)^{1-\frac{1}{q}}\left(\int_0^1\left|t^{\alpha}-\lambda\right|\left|f'\left(tx+(1-t)\,a\right)\right|^q dt\right)^{\frac{1}{q}} \\
(2.5) \quad & +\frac{(b-x)^{\alpha+1}}{b-a}\left(\int_0^1\left|t^{\alpha}-\lambda\right|dt\right)^{1-\frac{1}{q}}\left(\int_0^1\left|t^{\alpha}-\lambda\right|\left|f'\left(tx+(1-t)\,b\right)\right|^q dt\right)^{\frac{1}{q}}
\end{aligned}$$

Since$|f'|^q$ is $s-$convex on $[a,b]$ we get

$$\begin{aligned}\int_0^1 |t^{\alpha}-\lambda|\left|f'\left(tx+(1-t)a\right)\right|^q dt &\leq \int_0^1 |t^{\alpha}-\lambda|\left(t^s\left|f'(x)\right|^q+(1-t)^s\left|f'(a)\right|^q\right)dt \\ &= \left|f'(x)\right|^q A_2(\alpha,\lambda,s)+\left|f'(a)\right|^q A_3(\alpha,\lambda,s),\end{aligned} \tag{2.6}$$

$$\begin{aligned}\int_0^1 |t^{\alpha}-\lambda|\left|f'\left(tx+(1-t)b\right)\right|^q dt &\leq \int_0^1 |t^{\alpha}-\lambda|\left(t^s\left|f'(x)\right|^q+(1-t)^s\left|f'(b)\right|^q\right)dt \\ &= \left|f'(x)\right|^q A_2(\alpha,\lambda,s)+\left|f'(b)\right|^q A_3(\alpha,\lambda,s),\end{aligned} \tag{2.7}$$

where we use the fact that

$$\begin{aligned}\int_0^1 |t^{\alpha}-\lambda|(1-t)^s dt &= \int_0^{\lambda^{\frac{1}{\alpha}}}(\lambda-t^{\alpha})(1-t)^s dt+\int_{\lambda^{\frac{1}{\alpha}}}^1 (t^{\alpha}-\lambda)(1-t)^s dt \\ &= \lambda\int_0^{\lambda^{\frac{1}{\alpha}}}(1-t)^s dt-\int_0^{\lambda^{\frac{1}{\alpha}}}t^{\alpha}(1-t)^s dt+\int_{\lambda^{\frac{1}{\alpha}}}^1 t^{\alpha}(1-t)^s dt \\ &\quad -\lambda\int_{\lambda^{\frac{1}{\alpha}}}^1 (1-t)^s dt \\ &= \lambda\left[\frac{1-2\left(1-\lambda^{\frac{1}{\alpha}}\right)^{s+1}}{s+1}\right]+\int_0^1 t^{\alpha}(1-t)^s dt-2\int_0^{\lambda^{\frac{1}{\alpha}}}t^{\alpha}(1-t)^s dt \\ &= \lambda\left[\frac{1-2\left(1-\lambda^{\frac{1}{\alpha}}\right)^{s+1}}{s+1}\right]+\beta(\alpha+1,s+1)-2\beta\left(\lambda^{\frac{1}{\alpha}};\alpha+1,s+1\right),\end{aligned}$$

$$\int_0^1 |t^{\alpha}-\lambda|\,t^s dt=\frac{2\alpha\lambda^{1+\frac{s+1}{\alpha}}+s+1}{(s+1)(\alpha+s+1)}-\frac{\lambda}{s+1},$$

and by simple computation

$$\begin{aligned}\int_0^1 |t^{\alpha}-\lambda|\,dt &= \int_0^{\lambda^{\frac{1}{\alpha}}}(\lambda-t^{\alpha})\,dt+\int_{\lambda^{\frac{1}{\alpha}}}^1 (t^{\alpha}-\lambda)\,dt \\ &= \frac{2\alpha\lambda^{1+\frac{1}{\alpha}}+1}{\alpha+1}-\lambda.\end{aligned} \tag{2.8}$$

Hence, If we use (2.6), (2.7) and (2.8) in (2.5), we obtain the desired result. This completes the proof. □

Corollary 1. *Under the assumptions of Theorem 4 with* $q=1$, *the inequality (2.4) reduced to the following inequality*

$$\begin{aligned} |S_f(x,\lambda,\alpha,a,b)| &\leq \left\{ \frac{(x-a)^{\alpha+1}}{b-a} \left(|f'(x)| A_2(\alpha,\lambda,s) + |f'(a)| A_3(\alpha,\lambda,s) \right) \right. \\ & \left. + \frac{(b-x)^{\alpha+1}}{b-a} \left(|f'(x)| A_2(\alpha,\lambda,s) + |f'(b)| A_3(\alpha,\lambda,s) \right) \right\}. \end{aligned}$$

Corollary 2. *Under the assumptions of Theorem 4 with* $x=\frac{a+b}{2}$, $\lambda=\frac{1}{3}$,*from the inequality (2.4) we get the following Simpson type inequality for fractional integrals*

$$\begin{aligned} & \left| \frac{2^{\alpha-1}}{(b-a)^{\alpha-1}} S_f\left(\frac{a+b}{2},\frac{1}{3},\alpha,a,b\right) \right| \\ = & \left| \frac{1}{6}\left[f(a) + 4f\left(\frac{a+b}{2}\right) + f(b) \right] - \frac{\Gamma(\alpha+1)\,2^{\alpha-1}}{(b-a)^{\alpha}} \left[J^{\alpha}_{\left(\frac{a+b}{2}\right)^-} f(a) + J^{\alpha}_{\left(\frac{a+b}{2}\right)^+} f(b) \right] \right| \\ \leq & \frac{b-a}{4} A_1^{1-\frac{1}{q}}\left(\alpha,\frac{1}{3}\right) \left\{ \left(\left| f'\left(\frac{a+b}{2}\right) \right|^q A_2\left(\alpha,\frac{1}{3},s\right) + |f'(a)|^q A_3\left(\alpha,\frac{1}{3},s\right) \right)^{\frac{1}{q}} \right. \\ & \left. + \left(\left| f'\left(\frac{a+b}{2}\right) \right|^q A_2\left(\alpha,\frac{1}{3},s\right) + |f'(b)|^q A_3\left(\alpha,\frac{1}{3},s\right) \right)^{\frac{1}{q}} \right\}. \end{aligned}$$

Corollary 3. *Under the assumptions of Theorem 4 with* $x=\frac{a+b}{2}$, $\lambda=0$,*from the inequality (2.4) we get the following midpoint type inequality for fractional integrals*

$$\begin{aligned} & \left| \frac{2^{\alpha-1}}{(b-a)^{\alpha-1}} S_f\left(\frac{a+b}{2},0,\alpha,a,b\right) \right| \\ = & \left| f\left(\frac{a+b}{2}\right) - \frac{\Gamma(\alpha+1)\,2^{\alpha-1}}{(b-a)^{\alpha}} \left[J^{\alpha}_{\left(\frac{a+b}{2}\right)^-} f(a) + J^{\alpha}_{\left(\frac{a+b}{2}\right)^+} f(b) \right] \right| \\ \leq & \frac{b-a}{4}\left(\frac{1}{\alpha+1}\right)^{1-\frac{1}{q}} \left\{ \left[\frac{\left|f'\left(\frac{a+b}{2}\right)\right|^q}{\alpha+s+1} + |f'(a)|^q \beta(\alpha+1,s+1) \right]^{\frac{1}{q}} \right. \\ & \left. + \left[\frac{\left|f'\left(\frac{a+b}{2}\right)\right|^q}{\alpha+s+1} + |f'(b)|^q \beta(\alpha+1,s+1) \right]^{\frac{1}{q}} \right\}. \end{aligned}$$

Corollary 4. *Under the assumptions of Theorem 4 with* $\lambda=1$,*from the inequality (2.4) we get the following trapezoid type inequality for fractional integrals*

$$\begin{aligned} & \left| S_f\left(\frac{a+b}{2},1,\alpha,a,b\right) \right| \\ = & \left| \frac{(x-a)^{\alpha} f(a) + (b-x)^{\alpha} f(b)}{b-a} - \frac{\Gamma(\alpha+1)}{b-a} \left[J^{\alpha}_{x^-} f(a) + J^{\alpha}_{x^+} f(b) \right] \right| \\ \leq & \left(\frac{\alpha}{\alpha+1}\right)^{1-\frac{1}{q}} \left\{ \frac{(x-a)^{\alpha+1}}{b-a} \left[\frac{\alpha\,|f'(x)|^q}{(s+1)(\alpha+s+1)} + |f'(a)|^q \left(\frac{1}{s+1} - \beta(\alpha+1,s+1) \right) \right]^{\frac{1}{q}} \right. \\ & \left. + \frac{(b-x)^{\alpha+1}}{b-a} \left[\frac{\alpha\,|f'(x)|^q}{(s+1)(\alpha+s+1)} + |f'(b)|^q \left(\frac{1}{s+1} - \beta(\alpha+1,s+1) \right) \right]^{\frac{1}{q}} \right\} \end{aligned}$$

which is the same of the inequality in [8, Theorem 9].

Remark 1. *In Corollary 4, if we choose* $\alpha = 1$, *we get the same inequality in* [2, Theorem 7].

Corollary 5. *Let the assumptions of Theorem 4 hold. If* $|f'(x)| \leq M$ *for all* $x \in [a,b]$ *and* $\lambda = 0$, *then from the inequality (2.4) we get the following Ostrowski type inequality for fractional integrals*

$$\begin{aligned} &\left| \left[\frac{(x-a)^{\alpha} + (b-x)^{\alpha}}{b-a} \right] f(x) - \frac{\Gamma(\alpha+1)}{b-a} \left[J_{x^-}^{\alpha} f(a) + J_{x^+}^{\alpha} f(b) \right] \right| \\ \leq\ & M \left(\frac{1}{\alpha+1} \right)^{1-\frac{1}{q}} \left[\frac{1}{\alpha+s+1} + \beta(\alpha+1, s+1) \right]^{\frac{1}{q}} \left[\frac{(x-a)^{\alpha+1} + (b-x)^{\alpha+1}}{b-a} \right] \end{aligned}$$

for each $x \in [a,b]$.

Remark 2. *In Corollary 5, if we choose* $\alpha = 1$, *we get the same inequality in* [1, Theorem 4].

2.2. **For quasi-convex functions.**

Theorem 5. *Let* $f : I \subset [0,\infty) \to \mathbb{R}$ *be a differentiable function on* I° *such that* $f' \in L[a,b]$, *where* $a, b \in I^{\circ}$ *with* $a < b$. *If* $|f'|^q$ *is quasi-convex on* $[a,b]$ *for some fixed* $q \geq 1$, $x \in [a,b]$, $\lambda \in [0,1]$ *then the following inequality for fractional integrals holds*

$$\begin{aligned} (2.9)\ |S_f(x,\lambda,\alpha,a,b)| \ \leq\ & A_1(\alpha,\lambda) \left\{ \frac{(x-a)^{\alpha+1}}{b-a} \left(\sup\left\{ |f'(x)|^q, |f'(a)|^q \right\} \right)^{\frac{1}{q}} \right. \\ & \left. + \frac{(b-x)^{\alpha+1}}{b-a} \left(\sup\left\{ |f'(x)|^q, |f'(b)|^q \right\} \right)^{\frac{1}{q}} \right\} \end{aligned}$$

where

$$A_1(\alpha,\lambda) = \frac{2\alpha\lambda^{1+\frac{1}{\alpha}} + 1}{\alpha+1} - \lambda.$$

Proof. We proceed similarly as in the proof Theorem 4. Since $|f'|^q$ is quasi-convex on $[a,b]$, for all $t \in [0,1]$

$$|f'(tx + (1-t)a)|^q \leq \sup\left\{ |f'(x)|^q, |f'(a)|^q \right\}$$

and

$$|f'(tx + (1-t)b)|^q \leq \sup\left\{ |f'(x)|^q, |f'(b)|^q \right\}.$$

Hence, from the inequality (2.5) we get

$$\begin{aligned}
&\left|S_f\left(x,\lambda,\alpha,a,b\right)\right| \\
\leq\;& \left(\int_0^1 \left|t^{\alpha}-\lambda\right| dt\right)\left\{\frac{(x-a)^{\alpha+1}}{b-a}\left(\sup\left\{\left|f'(x)\right|^q,\left|f'(a)\right|^q\right\}\right)^{\frac{1}{q}}\right. \\
&\left.+\frac{(b-x)^{\alpha+1}}{b-a}\left(\sup\left\{\left|f'(x)\right|^q,\left|f'(b)\right|^q\right\}\right)^{\frac{1}{q}}\right\} \\
\leq\;& A_1\left(\alpha,\lambda\right)\left\{\frac{(x-a)^{\alpha+1}}{b-a}\left(\sup\left\{\left|f'(x)\right|^q,\left|f'(a)\right|^q\right\}\right)^{\frac{1}{q}}\right. \\
&\left.+\frac{(b-x)^{\alpha+1}}{b-a}\left(\sup\left\{\left|f'(x)\right|^q,\left|f'(b)\right|^q\right\}\right)^{\frac{1}{q}}\right\}
\end{aligned}$$

which completes the proof. □

Corollary 6. *Under the assumptions of Theorem 5 with* $q=1$, *the inequality (2.9) reduced to the following inequality*

$$\begin{aligned}
\left|S_f\left(x,\lambda,\alpha,a,b\right)\right| \leq\;& A_1\left(\alpha,\lambda\right)\left\{\frac{(x-a)^{\alpha+1}}{b-a}\left(\sup\left\{\left|f'(x)\right|,\left|f'(a)\right|\right\}\right)\right. \\
&\left.+\frac{(b-x)^{\alpha+1}}{b-a}\left(\sup\left\{\left|f'(x)\right|,\left|f'(b)\right|\right\}\right)\right\}.
\end{aligned}$$

Corollary 7. *Under the assumptions of Theorem 5 with* $x=\frac{a+b}{2}$, $\lambda=\frac{1}{3}$,*from the inequality (2.9) we get the following Simpson type inequality or fractional integrals*

$$\begin{aligned}
&\left|\frac{1}{6}\left[f(a)+4f\left(\frac{a+b}{2}\right)+f(b)\right]-\frac{\Gamma\left(\alpha+1\right)2^{\alpha-1}}{(b-a)^{\alpha}}\left[J^{\alpha}_{\left(\frac{a+b}{2}\right)^-}f(a)+J^{\alpha}_{\left(\frac{a+b}{2}\right)^+}f(b)\right]\right| \\
\leq\;& \frac{b-a}{4}A_1\left(\alpha,\frac{1}{3}\right)\left[\sup\left\{\left|f'\left(\frac{a+b}{2}\right)\right|,\left|f'(a)\right|\right\}+\sup\left\{\left|f'\left(\frac{a+b}{2}\right)\right|,\left|f'(b)\right|\right\}\right].
\end{aligned}$$

Corollary 8. *Under the assumptions of Theorem 5 with* $x=\frac{a+b}{2}$, $\lambda=0$,*from the inequality (2.9) we get the following midpoint type inequality or fractional integrals*

$$\begin{aligned}
&\left|f\left(\frac{a+b}{2}\right)-\frac{\Gamma\left(\alpha+1\right)2^{\alpha-1}}{(b-a)^{\alpha}}\left[J^{\alpha}_{\left(\frac{a+b}{2}\right)^-}f(a)+J^{\alpha}_{\left(\frac{a+b}{2}\right)^+}f(b)\right]\right| \\
\leq\;& \frac{b-a}{4}\left(\frac{1}{\alpha+1}\right)\left\{\left[\sup\left\{\left|f'\left(\frac{a+b}{2}\right)\right|^q,\left|f'(a)\right|^q\right\}\right]^{\frac{1}{q}}\right. \\
&\left.+\left[\sup\left\{\left|f'\left(\frac{a+b}{2}\right)\right|^q,\left|f'(b)\right|^q\right\}\right]^{\frac{1}{q}}\right\}.
\end{aligned}$$

Corollary 9. *Under the assumptions of Theorem 5 with* $\lambda = 1$,*from the inequality (2.9) we get the following trapezoid type inequality or fractional integrals*

$$\begin{aligned}
&\left|\frac{(x-a)^{\alpha} f(a) + (b-x)^{\alpha} f(b)}{b-a} - \frac{\Gamma(\alpha+1)}{b-a}\left[J_{x^-}^{\alpha} f(a) + J_{x^+}^{\alpha} f(b)\right]\right| \\
\leq \;& \left(\frac{\alpha}{\alpha+1}\right)\left\{\frac{(x-a)^{\alpha+1}}{b-a}\left[\sup\left\{\left|f'(x)\right|^{q}, \left|f'(a)\right|^{q}\right\}\right]^{\frac{1}{q}}\right. \\
&\left. + \frac{(b-x)^{\alpha+1}}{b-a}\left[\sup\left\{\left|f'(x)\right|^{q}, \left|f'(b)\right|^{q}\right\}\right]^{\frac{1}{q}}\right\}.
\end{aligned}$$

Corollary 10. *Let the assumptions of Theorem 5 hold. If* $|f'(x)| \leq M$ *for all* $x \in [a,b]$ *and* $\lambda = 0$, *then from the inequality (2.9) we get the following Ostrowski type inequality or fractional integrals*

$$\begin{aligned}
&\left|\left[\frac{(x-a)^{\alpha} + (b-x)^{\alpha}}{b-a}\right] f(x) - \frac{\Gamma(\alpha+1)}{b-a}\left[J_{x^-}^{\alpha} f(a) + J_{x^+}^{\alpha} f(b)\right]\right| \\
\leq \;& \frac{M}{\alpha+1}\left[\frac{(x-a)^{\alpha+1} + (b-x)^{\alpha+1}}{b-a}\right],
\end{aligned}$$

2.3. **For** $m-$**convex functions.** Similarly lemma 1, we can proved the following lemma

Lemma 2. *,Let* $f : I \subset \mathbb{R} \to \mathbb{R}$ *be a differentiable mapping on* I° *such that* $f' \in L[ma, mb]$, *where* $m \in (0,1]$, $ma, mb \in I$ *with* $a < b$. *Then for all* $x \in [a,b]$, $\lambda \in [0,1]$ *and* $\alpha > 0$ *we have:*

$$\begin{aligned}
S_f(mx, \lambda, \alpha, ma, mb) &= \frac{m^{\alpha}(x-a)^{\alpha+1}}{b-a}\int_0^1 \left(t^{\alpha} - \lambda\right) f'\left(tmx + m(1-t)a\right) dt \\
&+ \frac{m^{\alpha}(b-x)^{\alpha+1}}{b-a}\int_0^1 \left(\lambda - t^{\alpha}\right) f'\left(tmx + m(1-t)b\right) dt.
\end{aligned}$$

Theorem 6. *Let* $f : I \subset [0,\infty) \to \mathbb{R}$ *be a differentiable function on* I° *such that* $f' \in L[ma, mb]$, *where* $m \in (0,1]$, $a, b \in I^{\circ}$ *with* $a < b$. *If* $|f'|^q$ *is* $m-$*convex on* $[ma, mb]$ *for some fixed* $q \geq 1$, $x \in [a,b]$, $\lambda \in [0,1]$ *and* $\alpha > 0$ *then the following inequality for fractional integrals holds*

$$\begin{aligned}
&\left|S_f(mx, \lambda, \alpha, ma, mb)\right| \\
\leq \;& A_1(\alpha,\lambda)^{1-\frac{1}{q}}\left\{\frac{m^{\alpha}(x-a)^{\alpha+1}}{b-a}\left(\left|f'(mx)\right|^{q} A_2(\alpha,\lambda) + m\left|f'(a)\right|^{q} A_3(\alpha,\lambda)\right)^{\frac{1}{q}}\right. \\
&\left. + \frac{m^{\alpha}(b-x)^{\alpha+1}}{b-a}\left(\left|f'(mx)\right|^{q} A_2(\alpha,\lambda) + m\left|f'(b)\right|^{q} A_3(\alpha,\lambda)\right)^{\frac{1}{q}}\right\}
\end{aligned} \tag{2.10}$$

where

$$\begin{aligned}
A_1(\alpha,\lambda) &= \frac{2\alpha\lambda^{1+\frac{1}{\alpha}}+1}{\alpha+1}-\lambda, \\
A_2(\alpha,\lambda) &= \frac{\alpha\lambda^{1+\frac{2}{\alpha}}+1}{\alpha+2}-\frac{\lambda}{2}, \\
A_3(\alpha,\lambda) &= \frac{2\alpha\lambda^{1+\frac{1}{\alpha}}+1}{\alpha+1}-\frac{\alpha\lambda^{1+\frac{2}{\alpha}}+1}{\alpha+2}-\frac{\lambda}{2}.
\end{aligned}$$

Proof. We proceed similarly as in the proof Theorem 4. From Lemma 2, property of the modulus and using the power-mean inequality we have

$$\begin{aligned}
&\left|S_f(mx,\lambda,\alpha,ma,mb)\right| \\
\leq\ & \frac{m^\alpha (x-a)^{\alpha+1}}{b-a}\left(\int_0^1 |t^\alpha-\lambda|\,dt\right)^{1-\frac{1}{q}}\left(\int_0^1 |t^\alpha-\lambda|\left|f'(tmx+m(1-t)a)\right|^q dt\right)^{\frac{1}{q}} \\
& \frac{m^\alpha (b-x)^{\alpha+1}}{b-a}\left(\int_0^1 |t^\alpha-\lambda|\,dt\right)^{1-\frac{1}{q}}\left(\int_0^1 |t^\alpha-\lambda|\left|f'(tmx+m(1-t)b)\right|^q dt\right)^{\frac{1}{q}}. \qquad (2.11)
\end{aligned}$$

Since $|f'|^q$ is $m-$convex on $[ma,mb]$, for all $t\in[0,1]$

$$\left|f'(tmx+m(1-t)a)\right|^q \leq t\left|f'(mx)\right|^q+m(1-t)\left|f'(a)\right|^q$$

and

$$\left|f'(tmx+m(1-t)b)\right|^q \leq t\left|f'(mx)\right|^q+m(1-t)\left|f'(b)\right|^q.$$

Hence by simple computation we get

$$\begin{aligned}
\int_0^1 |t^\alpha-\lambda|\left|f'(tmx+m(1-t)a)\right|^q dt &\leq \int_0^1 |t^\alpha-\lambda|\,t\left|f'(mx)\right|^q+m(1-t)\left|f'(a)\right|^q dt \\
&= \left|f'(mx)\right|^q\left(\frac{\alpha\lambda^{1+\frac{2}{\alpha}}+1}{\alpha+2}-\frac{\lambda}{2}\right) \\
&\quad +m\left|f'(a)\right|^q\left(\frac{2\alpha\lambda^{1+\frac{1}{\alpha}}+1}{\alpha+1}-\frac{\alpha\lambda^{1+\frac{2}{\alpha}}+1}{\alpha+2}-\frac{\lambda}{2}\right), \qquad (2.12)
\end{aligned}$$

and similarly

$$\begin{aligned}
\int_0^1 |t^\alpha-\lambda|\left|f'(tmx+m(1-t)b)\right|^q dt &\leq \int_0^1 |t^\alpha-\lambda|\,t\left|f'(mx)\right|^q+m(1-t)\left|f'(b)\right|^q dt \\
&= \left|f'(mx)\right|^q\left(\frac{\alpha\lambda^{1+\frac{2}{\alpha}}+1}{\alpha+2}-\frac{\lambda}{2}\right) \\
&\quad +m\left|f'(b)\right|^q\left(\frac{2\alpha\lambda^{1+\frac{1}{\alpha}}+1}{\alpha+1}-\frac{\alpha\lambda^{1+\frac{2}{\alpha}}+1}{\alpha+2}-\frac{\lambda}{2}\right). \qquad (2.13)
\end{aligned}$$

If we use (2.12), (2.13) and (2.8) in (2.11), we obtain the desired result. This completes the proof. □

Corollary 11. *Under the assumptions of Theorem 6 with* $q = 1$*, the inequality (2.10) reduced to the following inequality*

$$\begin{aligned} \left|S_f\left(mx,\lambda,\alpha,ma,mb\right)\right| \leq & \left\{\frac{m^{\alpha}\left(x-a\right)^{\alpha+1}}{b-a}\left(\left|f'\left(mx\right)\right|A_2\left(\alpha,\lambda\right)+m\left|f'\left(a\right)\right|A_3\left(\alpha,\lambda\right)\right)\right. \\ & \left.+\frac{m^{\alpha}\left(b-x\right)^{\alpha+1}}{b-a}\left(\left|f'\left(mx\right)\right|A_2\left(\alpha,\lambda\right)+m\left|f'\left(b\right)\right|A_3\left(\alpha,\lambda\right)\right)\right\}. \end{aligned}$$

Corollary 12. *Under the assumptions of Theorem 6 with* $x = \frac{a+b}{2}$, $\lambda = \frac{1}{3}$*,from the inequality (2.10) we get the following Simpson type inequality or fractional integrals*

$$\begin{aligned} & \left|\frac{2^{\alpha-1}}{m^{\alpha}\left(b-a\right)^{\alpha-1}}S_f\left(m\left(\frac{a+b}{2}\right),\frac{1}{3},\alpha,ma,mb\right)\right| \\ = & \left|\frac{1}{6}\left[f(ma)+4f\left(\frac{m\left(a+b\right)}{2}\right)+f(mb)\right]-\frac{\Gamma\left(\alpha+1\right)2^{\alpha-1}}{m^{\alpha}\left(b-a\right)^{\alpha}}\left[J^{\alpha}_{\left(\frac{m(a+b)}{2}\right)^-}f(ma)+J^{\alpha}_{\left(\frac{m(a+b)}{2}\right)^+}f(mb)\right]\right| \\ \leq & \frac{m\left(b-a\right)}{4}A_1^{1-\frac{1}{q}}\left(\alpha,\frac{1}{3}\right)\left\{\left(\left|f'\left(\frac{m\left(a+b\right)}{2}\right)\right|^q A_2\left(\alpha,\frac{1}{3}\right)+m\left|f'\left(a\right)\right|^q A_3\left(\alpha,\frac{1}{3}\right)\right)^{\frac{1}{q}}\right. \\ & \left.+\left(\left|f'\left(\frac{m\left(a+b\right)}{2}\right)\right|^q A_2\left(\alpha,\frac{1}{3}\right)+m\left|f'\left(b\right)\right|^q A_3\left(\alpha,\frac{1}{3}\right)\right)^{\frac{1}{q}}\right\}. \end{aligned}$$

Corollary 13. *Under the assumptions of Theorem 6 with* $x = \frac{a+b}{2}$, $\lambda = 0$*,from the inequality (2.10) we get the following midpoint type inequality or fractional integrals*

$$\begin{aligned} & \frac{2^{\alpha-1}}{m^{\alpha}\left(b-a\right)^{\alpha-1}}\left|S_f\left(m\left(\frac{a+b}{2}\right),0,\alpha,ma,mb\right)\right| \\ = & \left|f\left(\frac{m\left(a+b\right)}{2}\right)-\frac{\Gamma\left(\alpha+1\right)2^{\alpha-1}}{m^{\alpha}\left(b-a\right)^{\alpha}}\left[J^{\alpha}_{\left(\frac{m(a+b)}{2}\right)^-}f(ma)+J^{\alpha}_{\left(\frac{m(a+b)}{2}\right)^+}f(mb)\right]\right| \\ \leq & \frac{m\left(b-a\right)}{4}\left(\frac{1}{\alpha+1}\right)\left(\frac{1}{\alpha+2}\right)^{\frac{1}{q}}\left\{\left[\left(\alpha+1\right)\left|f'\left(\frac{m\left(a+b\right)}{2}\right)\right|^q+m\left|f'\left(a\right)\right|^q\right]^{\frac{1}{q}}\right. \\ & \left.+\left[\left(\alpha+1\right)\left|f'\left(\frac{m\left(a+b\right)}{2}\right)\right|^q+m\left|f'\left(b\right)\right|^q\right]^{\frac{1}{q}}\right\}. \end{aligned}$$

Corollary 14. *Under the assumptions of Theorem 6 with* $\lambda = 1$*,from the inequality (2.10) we get the following trapezoid type inequality or fractional integrals*

$$\begin{aligned} & \left|S_f\left(mx,1,\alpha,ma,mb\right)\right| \\ = & \left|\frac{\left(x-a\right)^{\alpha}f(ma)+\left(b-x\right)^{\alpha}f(mb)}{b-a}-\frac{\Gamma\left(\alpha+1\right)}{m^{\alpha+1}\left(b-a\right)}\left[J^{\alpha}_{mx^-}f(ma)+J^{\alpha}_{mx^+}f(mb)\right]\right| \\ \leq & \left(\frac{\alpha}{\alpha+1}\right)\left(\frac{1}{2\left(\alpha+2\right)}\right)^{\frac{1}{q}}\left\{\frac{\left(x-a\right)^{\alpha+1}}{b-a}\left[\left(\alpha+1\right)\left|f'\left(mx\right)\right|^q+\left(\alpha+3\right)m\left|f'\left(a\right)\right|\right]^{\frac{1}{q}}\right. \\ & \left.+\frac{\left(b-x\right)^{\alpha+1}}{b-a}\left[\left(\alpha+1\right)\left|f'\left(mx\right)\right|^q+\left(\alpha+3\right)m\left|f'\left(b\right)\right|\right]^{\frac{1}{q}}\right\}. \end{aligned}$$

Corollary 15. *Let the assumptions of Theorem 6 hold. If* $|f'(u)| \leq M$ *for all* $u \in [ma, mb]$ *and* $\lambda = 0$*, then from the inequality (2.10) we get the following*

Ostrowski type inequality or fractional integrals

$$\left|\left[\frac{(x-a)^{\alpha}+(b-x)^{\alpha}}{b-a}\right]f(mx)-\frac{\Gamma(\alpha+1)}{m^{\alpha+1}(b-a)}\left[J_{mx^{-}}^{\alpha}f(ma)+J_{mx^{+}}^{\alpha}f(mb)\right]\right|$$
$$\leq \quad mM\left(\frac{1}{\alpha+1}\right)\left(\frac{\alpha+m+1}{\alpha+2}\right)^{\frac{1}{q}}\left[\frac{(x-a)^{\alpha+1}+(b-x)^{\alpha+1}}{b-a}\right],$$

for each $x\in[a,b]$.

References

[1] M. Alomaria, M. Darus, S.S. Dragomir, P. Cerone, Ostrowski type inequalities for functions whose derivatives ares-convex in the second sense, Applied Mathematics Letters 23 (2010) 1071–1076.

[2] M. Avci, H. Kavurmaci and M.E. Ozdemir, New inequalities of Hermite-Hadamard type via $s-$convex functions in the second sense with applications, Appl. Math. Comput., 217 (2011) 5171–5176.

[3] Z. Dahmani, On Minkowski and Hermite-Hadamard integral inequalities via fractional via fractional integration, Ann. Funct. Anal. 1 (1) (2010), 51-58

[4] R. Gorenflo, F. Mainardi, Fractional calculus; integral and differential equations of fractional order, Springer Verlag, Wien (1997), 223-276.

[5] H. Hudzik and L. Maligranda, Some remarks on $s-$convex functions, Aequationes Math. 48 (1994), 100-111.

[6] D.A. Ion, Some estimates onthe Hermite-Hadamard inequality through quasi-convex functions, Annals of University of Craiova, Math. Comp. Sci. Ser. 34 (2007) 82-87.

[7] S. Miller and B. Ross, An introduction to the Fractional Calculus and Fractional Differential Equations, John Wiley & Sons, USA (1993), 2.

[8] M.E. Ozdemir, M. Avci, H. Kavurmaci, Hermite-Hadamard type inequalities for $s-$convex and $s-$concave functions via fractional integrals, arXiv:1202.0380v1.

[9] I. Podlubni, Fractional Differential Equations, Academic Press, San Diego, 1999.

[10] J. Pečarić, F. Proschan, and Y.L. Tong, Convex Functions, Partial Orderings and Statistical Applications, Academic Press Inc. 1992.

[11] E. Set, New inequalities of Ostrowski type for mapping whose derivatives are s-convex in the second sense via fractional integrals, Computers and Math. with Appl. 63 (2012) 1147-1154.

[12] M.Z. Sarıkaya, E. Set, H. Yaldız and N. Başak, Hermite-Hadamard's inequalities for fractional integrals and related fractional inequalities, Mathematical and Computer Modelling, DOI:10.1016/j.mcm.2011.12.048.

[13] G. Toader, Some generalizations ofthe convexity, Proceedings of The Colloquium On Approximation And Optimization, Univ. Cluj-Napoca, Cluj-Napoca,1985, 329-338.

Department of Mathematics, Faculty of Sciences and Arts, Giresun University, Giresun, Turkey

E-mail address: `imdat.iscan@giresun.edu.tr`